

# A MULTILINEAR PROOF OF JACOBI'S FORMULA

Jules Jacobs

August 25, 2022

Jacobi's formula gives the determinant of the matrix exponential in terms of the trace:

$$\det(e^{tA}) = e^{t \operatorname{tr}(A)}$$

The usual proof of this uses Laplace expansion of the determinant. Here is a proof that relies on multilinearity of the determinant:

Consider the function  $f(t) = \det(e^{tA})$ . To show that  $f(t) = e^{t \operatorname{tr}(A)}$  it suffices to show that  $f$  satisfies the differential equation that defines  $e^{t \operatorname{tr}(A)}$ :

$$\begin{aligned} f(0) &= 1 \\ f'(t) &= \operatorname{tr}(A)f(t) \end{aligned}$$

The condition  $f(0) = 1$  follows immediately, so it remains to show  $f'(t)/f(t) = \operatorname{tr}(A)$ . In other words, we have to show that  $\det(e^{tA})' \det(e^{-tA}) = \operatorname{tr} A$ .

To calculate the derivative of a determinant, first consider the derivative of a product:

$$(a_1 \cdot a_2 \cdots a_n)' = (a'_1 \cdot a_2 \cdots a_n) + (a_1 \cdot a'_2 \cdots a_n) + \cdots + (a_1 \cdot a_2 \cdots a'_n)$$

The analogous derivative formula works for any multilinear function, such as the determinant, where  $\det(a_1 | a_2 | \cdots | a_n)$  is the determinant of the matrix with columns  $a_1, a_2, \dots, a_n$ :

$$\det(a_1 | a_2 | \cdots | a_n)' = \det(a'_1 | a_2 | \cdots | a_n) + \det(a_1 | a'_2 | \cdots | a_n) + \cdots + \det(a_1 | a_2 | \cdots | a'_n)$$

Using this derivative formula and  $(e^{tA})' = Ae^{tA}$  and  $e^{tA}e^{-tA} = I$ , we get:

$$\det(e^{tA})' \det(e^{-tA}) = \det(A_1 | I_2 | \cdots | I_n) + \det(I_1 | A_2 | \cdots | I_n) + \cdots + \det(I_1 | I_2 | \cdots | A_n) = \operatorname{tr}(A)$$

This completes the proof.

A similar argument shows that  $\det(A)' = \operatorname{tr}(A^{-1}A')\det(A)$ .